

Constraints on Intervening Stellar Populations Toward the Large Magellanic Cloud

Dennis Zaritsky

UCO/Lick Observatories and Department of Astronomy and Astrophysics,
Univ. of California at Santa Cruz, Santa Cruz, CA, 95064
Electronic mail: dennis@ucolick.org

Stephen A. Shectman and Ian Thompson

Carnegie Observatories, 813 Santa Barbara St., Pasadena, CA 91101
Electronic mail: shec@ociw.edu, ian@ociw.edu

Jason Harris and D.N.C. Lin

Department of Astronomy and Astrophysics, Univ. of California at Santa Cruz,
Santa Cruz, CA, 95064
Electronic mail: jharris@ucolick.org, lin@ucolick.org

ABSTRACT

The suggestion by Zaritsky & Lin (1997; ZL) that a vertical extension of the red clump feature in color-magnitude diagrams (CMDs) of the Large Magellanic Cloud (LMC) is consistent with a significant population of foreground stars to the LMC that could account for the observed microlensing optical depth (Renault *et al.* 1997; Alcock *et al.* 1997a) has been challenged by various investigators (cf. Alcock *et al.* 1997b, Gallart 1998, Bennett 1998, Gould 1998, Beaulieu & Sackett 1998, and Ibata, Lewis, & Beaulieu 1998). We respond by (1) examining each of the challenges presented to determine whether any or all of those arguments invalidate the claims made by ZL and (2) presenting new photometric and spectroscopic data obtained in an attempt to resolve this issue. We systematically discuss why the objections raised so far do not unequivocally refute ZL's claim. We conclude that although the CMD data do not mandate the existence of a foreground population, they are entirely consistent with a foreground population associated with the LMC that contributes significantly ($\sim 50\%$) to the observed microlensing optical depth. From our new data, we conclude that $\lesssim 40\%$ of the VRC stars are young, massive red clump stars because (1) synthetic color-magnitude diagrams created using the star formation history derived independently from HST data (Geha *et al.* 1998) suggest that $< 50\%$ of the VRC stars are young, massive red clump stars, (2) the angular distribution of the VRC stars is more uniform than that of the young (age < 1 Gyr) main sequence stars, and (3) the velocity dispersion of the VRC stars in the region of the LMC examined by ZL, 18.4 ± 2.8 km sec $^{-1}$ (95% confidence limits), is inconsistent with the expectation for a young disk population. Each of these arguments is predicated on assumptions and the conclusions are uncertain. Therefore, an exact determination of the contribution to the microlensing optical depth by the various hypothesized foreground populations, and the subsequent conclusions regarding the existence of halo MACHOs, requires

a detailed knowledge of many complex astrophysical issues, such as the IMF, star formation history, and post-main sequence stellar evolution.

Subject headings: dark matter — Galaxy: halo — Magellanic Clouds

1. Introduction

The solution of the dark matter problem is a fundamental goal of current astronomical research. A particularly novel and ambitious approach involves the detection of gravitational lensing due to dark objects in the Galactic halo passing between us and background sources such as the stars in the Magellanic Clouds (Paczynski 1986). Several groups (EROS, cf. Aubourg *et al.* 1993; MACHO, cf. Alcock *et al.* 1997a; and OGLE, cf. Udalski, Kubiak, Szymanski 1997) have undertaken multi-year observing programs to detect such lensing events toward the Small and Large Magellanic Clouds (SMC and LMC). The fundamental successes of those surveys are that they convincingly identify microlensing events and that their derived microlensing optical depths are consistent. Using standard models for the stellar populations of the Galaxy and LMC, the observed microlensing optical depth, τ_μ , implies that $\sim 50\%$ of the Galactic halo dark matter out to the radius of the LMC is in the form of massive compact objects, MACHOs (Alcock *et al.* 1997a). The importance of this conclusion to our understanding of a wide range of astrophysical topics is manifest. Is this conclusion robust?

Several authors have by now questioned various assumptions leading to this conclusion. For example, Sahu (1994) proposed that LMC self-lensing (lensing of LMC stars by other LMC stars) is greater than in the standard model and can be sufficient to account for τ_μ ; Zhao (1998a) proposed that an associated intervening stellar population, such as a dwarf galaxy along the line of sight, might provide a significant fraction of the total optical depth; and Evans *et al.* (1998) proposed that the Galactic disk may be sufficiently warped and flared to provide $\sim 50\%$ of τ_μ . Zaritsky & Lin (1997; hereafter ZL) proposed that the vertical extension of the red clump feature (termed the VRC) observed in a million star color-magnitude diagram, CMD, of an LMC region observed as part of the Magellanic Clouds Photometric Survey (MCPS, Zaritsky, Harris & Thompson 1997; hereafter ZHT) is consistent with the existence of either an associated or unassociated intervening stellar population that is as much as 15 kpc closer to us than the LMC and that this population has sufficient mass density to account for a large fraction, and possibly all, of τ_μ .

To reach this conclusion, ZL subtracted a simple model of the red clump distribution and found a residual population that, if foreground, corresponds to stars ~ 15 kpc closer to us than the LMC. This model greatly simplified the analysis of the possible effect of an intervening population, but has led to some confusion in the literature. Because

the VRC is a continuous extension of the red clump toward brighter magnitudes, it may arise from stars distributed continuously between 0 and 15 kpc from the LMC. Therefore, the interpretation of the VRC impacts a variety of intervening-population scenarios, ranging from that of a detached stellar population unassociated with the LMC to that of a gravitationally bound halo or “thick” disk LMC population. The theoretical claims and possible observational evidence for intervening populations triggered a flurry of rebuttal papers (Alcock *et al.* 1997b, Gallart 1998, Bennett 1998, Gould 1998, Johnston 1998, Beaulieu & Sackett 1998, Ibata, Lewis, & Beaulieu 1998).

We systematically re-examine the possibility of a foreground population in two ways: (1) we test each of the arguments presented in the rebuttal papers to determine whether any or all of those arguments invalidate the most general claims made by ZL and (2) we present new photometric and spectroscopic data in an attempt to resolve this issue. We define our interpretation of Occam’s razor in §1.1. In §2 we describe our new data, in §3 we discuss the rebuttal papers and examine their conclusions, and in §4 we discuss the implications of the new data presented in §2. The discussion in §3 illustrates how difficult it is to eliminate the possibility of a foreground population responsible for the microlensing. In §4 we demonstrate that the nature of the VRC is complex and that stars in that region of the CMD must have a variety of origins. The myriad of uncertainties in the interpretation of the data and in the models applied to convert the observations into a microlensing optical depth must propagate into an uncertainty in any interpretation of the microlensing events. We conclude that due to the unresolved systematic uncertainties the microlensing data are yet unable to demonstrate the existence of halo MACHOs.

1.1. Occam’s Razor Revisited

Because we find microlensing by normal stars distributed in unexpected ways to be more plausible than microlensing by unknown objects distributed in a smooth halo, we assert that a convincing case for halo MACHOs can only be made after the sum of all plausible stellar populations is categorically eliminated as the source of τ_μ . Therefore, our approach is to aggressively attempt to reconcile the possibility of a foreground population that accounts for a significant fraction ($\sim 50\%$) of τ_μ with the various challenges raised by the rebuttal papers. If we can accommodate such a foreground population within parameter ranges allowed by observational or model uncertainties, then the interpretation of the microlensing optical depth is dominated by uncertainties in the parameters of the foreground population. Given the impact of the existence of halo MACHOs on a wide variety of disciplines, we believe that it is insufficient to argue that the MACHO interpretation is preferable to other interpretations that make moderate adjustments to such ill-determined quantities as the field initial mass function in the LMC or the star formation history of the LMC. We consider arguments

that present alternative interpretations for the VRC population or that conclude that the optical depth from such a population may be negligible ($\lesssim 10\%$), but which do not *convincingly eliminate* the presence or effect of such intervening populations on τ_μ , to be incomplete (even though they may be correct). We cede from the beginning that models exist for which the effect of the intervening populations are negligible — we address whether such models are unavoidable.

2. Data

2.1. Photometry

The photometric data come from an ongoing survey of the Magellanic Clouds (ZHT). We have now reduced data from an area in the LMC that is more than three times larger than that available to ZL. The new area also contains one of the most vigorous sites of recent (< 1 Gyr) star formation in the LMC, the Constellation III area (Shapley 1956), and so provides a range of stellar populations with different star formation histories and stellar densities. Photometric incompleteness becomes a serious factor only for $V > 21$ or in high density regions, such as the centers of stellar clusters (ZHT). Incompleteness is not a problem for this study of red clump or upper ($V < 19.5$) main sequence stars. The area being discussed, Figure 1, is centered at approximately $\alpha = 5^h20^m$ and $\delta = -66^\circ48'$ and has an irregular shape due to idiosyncrasies of the survey that do not affect our current analysis. The area discussed by ZL is roughly the lower right quarter of Figure 1. The stellar catalog consists of α, δ (2000.0) and $UBVI$ photometry and associated uncertainties. Stars must be detected in B and V to enter the catalog, but not necessarily in U and I . The current catalog contains ~ 2.5 million stars with $V < 21$ and 4 million in total.

2.2. Spectroscopy

The spectroscopic data were obtained using the Las Campanas 2.5m du Pont telescope and the multifiber spectrograph (Shectman *et al.* 1992) during the nights of January 19-23, 1998. This spectrograph obtains up to 128 spectra simultaneously. Due to the need to compromise between maximum spectral resolution and signal-to-noise, we used different observational setups for the brighter VRC stars than for the fainter red clump (RC) stars. We used a 1200 l/mm grating blazed at 5000Å in second order to observe the VRC stars (2.3 Å resolution) and in first order to observe the RC stars (3.8 Å resolution). The spectrograph uses a 2D-Frutti detector, which is blue sensitive, so we concentrate on the spectral region between 3800Å and 4800Å. Therefore, velocities are measured primarily using the Ca H and K lines.

One of our primary concerns for these observations is the contamination of spectra by other stars within the 3 arcsec fiber aperture. The fields in the LMC are crowded

and some stars, typically fainter than VRC or RC stars, are always present within a few arcsec of a random location. To quantify this problem and statistically remove “sky” from object spectra, we obtained spectra at a position offset by 8 arcsec from the target position. The offset frames were interleaved between target observations. The exposure times for individual exposures were 3000 sec for target and 1000 sec for sky. Several exposures, both target and sky, were taken for each field. The total exposure times were 11000 sec for VRC Field 1 ($\alpha = 5^h 6^m \delta = -67^\circ 42'$), 20000 sec for VRC Field 2 ($\alpha = 5^h 11^m \delta = -67^\circ 5'$), and 10000 for RC field 1 ($\alpha = 5^h 6^m \delta = -67^\circ 42'$). Each set of target and sky observations are bracketed by calibration spectra (internal hollow cathode lamp exposures). Dark frames and fiber throughput calibration frames (incandescent lamp exposures) were taken before each night’s observations.

The initial selection of targets was done using the color-magnitude diagrams originally used to identify the VRC (ZL). The VRC is defined using $3.1 < C < 3.4$ and $18 < V < 18.75$, where $C \equiv 0.565(B - I) + 0.825(U - V + 1.15)$. The quantity C was defined by ZL to remove a slight color-magnitude dependence of the red clump. In practice, it is only a minor modification and selecting the VRC in the $V - I, V$ space would produce a similar sample of stars. The RC region is defined by $3.1 < C < 3.4$ and $19 < V < 19.3$. The definitions for the two populations are more restrictive in color than those used by ZL and elsewhere in this paper to minimize contamination by other populations. We observe two non-overlapping regions on the sky located in the region of the LMC discussed by ZL. The stars were ranked in terms of isolation from other bright stars. During the otherwise random fiber assignment, the most isolated stars are given preference to minimize contamination.

The data reduction consists of the following steps: (1) normalize long dark frames to corresponding exposure time and subtract from all frames, (2) trace the incandescent lamp spectra (to define the apertures on the detector using a high S/N exposure) and extract it (to measure the relative throughputs of the fibers), (3) extract object and sky apertures using the apertures as defined by the incandescent lamp exposures only allowing a recentering of the reference apertures, (4) correct for fiber throughput differences by applying the correction factors measured from the incandescent lamp exposure, (5) derive wavelength solution for each fiber using calibration spectra obtained just before and after the science exposures, (6) apply wavelength solution to sky and object apertures, (7) combine sky spectra to form a single average sky spectrum for each sky position, (8) subtract sky spectrum from each object spectrum (after correcting for relative exposure times of sky and object), and (9) combine corresponding object spectra from multiple exposures of the same field taken during a night. The final product is a set of one-dimensional spectra of VRC and RC stars.

We measure velocities using standard cross-correlation software (XCSAO in IRAF). The template spectrum is constructed from the data themselves. We use a spectrum of a Galactic K giant to measure initial velocity estimates for the VRC stars. The spectra

of VRC stars with reliable correlation velocities ($R \geq 6$), are corrected back to zero radial velocity and combined to create a template with the same spectral characteristics of the sample stars (thereby avoiding template mismatch problems). We iterate this procedure until the number of reliable velocity measurements converges. The final template spectrum is shown in Figure 2. Because of this procedure, the mean velocity measured for the VRC stars is 0 km sec^{-1} . The RC stars could possibly have a mean velocity offset, and Galactic stars should have $v \sim -275 \text{ km sec}^{-1}$ (measurements of the LMC systematic velocity vary by only a few km sec^{-1} : 274 km sec^{-1} from H I observations of Luks & Rohlfs (1992) and 278 km sec^{-1} from a sample of planetary nebulae observed by Meatheringham *et al.* 1988).

It is critical to determine the observational uncertainties precisely because both the mean velocity difference between the VRC and RC stars and the velocity dispersions of the VRC and RC stars are expected to be comparable to the velocity uncertainty in any single spectrum ($\sim 10 \text{ km sec}^{-1}$). The cross correlation analysis calculates a measure of the uncertainty based on the strength of the correlation peak relative to the noise of the correlation function, but this calculation can underestimate the uncertainties — especially when there are systematic errors (eg. unstable wavelength solutions and poor sky subtraction). We conduct several tests to determine the reliability of the calculated uncertainties: (1) calibration spectra are cross-correlated against other calibration spectra to test the stability of the wavelength solution, (2) the velocities obtained for target stars reduced in the standard manner are compared with the velocities obtained when no sky spectra are subtracted to determine the maximum errors caused by poor sky subtraction, and (3) the velocities of VRC stars in Field 2 that were observed on two nights are compared.

The first test involves cross-correlating the self-calibrated calibration lamp spectra. Ideally, these would all have zero relative velocity, but in practice because of the sparse sampling of the wavelength solution, line centering errors, and polynomial fitting to the dispersion function, the spectra can appear to be shifted relative to one another. We have randomly selected one aperture from one exposure to act as a velocity template and cross correlated calibration lamps from various nights and all the apertures with this one template. The distribution of velocities has a mean of $-3.4 \pm 0.4 \text{ km sec}^{-1}$ and a dispersion of $2.6 \pm 0.3 \text{ km sec}^{-1}$. The distribution is Gaussian (see left panel Figure 3). The uncertainties introduced by errors in the wavelength calibration, especially because each stellar velocity consists of at least four different exposures (each with a different wavelength calibration) is negligible.

The second test is aimed at determining the uncertainties introduced by sky subtraction and contamination. We compare the results from one target field, with the results for the same stars when no sky subtraction is performed. This comparison should provide an upper limit to the velocity uncertainty associated with contamination and poor sky subtraction. As shown in the right panel of Figure 3, the distribution

of velocity differences is non-Gaussian and centered off zero (as expected, because the sky contributes a variable amplitude, but constant velocity signal to the spectrum). Because the best fit Gaussian has a dispersion of 5 km sec^{-1} , sky subtraction is not as serious a problem as was originally envisioned. The contribution to the overall dispersion, which should be less than in this extreme scenario, must be well less than 5 km sec^{-1} added in quadrature.

The third test involves comparing the measured velocities for the stars in the one target field that was observed on separate nights. We apply the criteria that $R \geq 6$ for an acceptable velocity, as we apply for all of our velocity measurements. We evaluate $D \equiv v_1 - v_2$, where v_1 and v_2 are the measured velocities from the first and second night respectively. The distribution of D should be Gaussian with mean zero and dispersion given by the standard error propagation, $\sigma^2 = \sigma_{v1}^2 + \sigma_{v2}^2$. The distribution of $D' \equiv D/\sigma$ should be Gaussian with mean zero and a dispersion of one if there are no systematic errors and the uncertainties calculated by the cross-correlation software are accurate. For the 80 stars that satisfy our correlation criteria in both night's data, the distribution is well fit by a Gaussian ($\chi^2 = 0.3$) with a mean of 0.0 ± 0.6 (indicating no significant nightly zero point shift) and a dispersion of $1.2^{+0.6}_{-0.3}$ (Figure 4). This result indicates that the uncertainties generated by the cross-correlation package are consistent with the scatter present in the velocities determined from different nights and that these uncertainties are reliable estimates of the true uncertainties. The data, including (α, δ) , V and B magnitudes, velocities, and velocity uncertainties, are available from the first author. The conclusions we draw from the data are discussed in §4.

3. Discussion

3.1. Previous Studies

A variety of studies, mostly in response to ZL, challenge the suggestion that foreground material traced by the VRC could account for τ_μ . The argument presented by ZL consisted of several steps, and each has been investigated in the series of papers discussed below. In this section, we discuss the studies by Alcock *et al.* (1997b), Gallart (1998), Bennett (1998), Gould (1998), Johnston (1998), and Beaulieu & Sackett (1998) in chronological order of publication. We discuss the results of Ibata, Lewis, & Beaulieu (1998) in §4 in the context of our radial velocity measurements.

3.1.1. Alcock *et al.* 1997b

The most direct test for the presence of intervening populations is to measure distances to stars (or lenses) along the line of sight. Alcock *et al.* use the MACHO database to search for variable stars for which distances can be derived (Cepheids

and RR Lyrae). They do not find a population of foreground Cepheids, but Cepheids are somewhat unlikely in an intervening population since their presence would imply very recent (age $\lesssim \text{few} \times 10^8$ yrs; Kippenhahn & Smith 1969; Efremov 1978; Grebel & Brandner 1998) star formation. RR Lyrae are more likely to be present in an intervening population and Zhao (1998a) noted that the data of Payne-Gaposchkin (1971) contained a tantalizing clump of RR Lyrae candidates at a distance of ~ 20 kpc.

Alcock *et al.* found no excess of RR Lyrae over the number expected in a flattened ($b/a = 0.6$) halo model that was normalized to produce the total number of stars identified to a distance of 40 kpc along the line of sight to the LMC. However, due to possible confusion with blends, LMC RR Lyrae, and other variable stars they imposed a selection cut of $V < 18$. For their adopted RR Lyrae absolute V magnitude of +0.4, this selection cut corresponds to a limiting distance of 33 kpc. Therefore, while their result argues against the Zhao hypothesis of an intervening galaxy at ~ 20 kpc, it does not place strong limits on possible stellar populations at distances of 35 to 50 kpc.

Alcock *et al.* also note that there are no tell-tale features of a foreground population visible in deep HST CMDs. Because of the expected large number of low mass stars for each “high” mass star (*e.g.*, red clump star), a foreground population may be more statistically distinct along the lower main sequence or main sequence turnoff region of a CMD than at brighter magnitudes. However, such comparisons, when the hypothesized foreground population is a small fraction of the total population, are difficult because of the nearly vertical distribution of main sequence stars in CMDs. In Figure 5 we compare an original HST CMD of the LMC (courtesy of J. Holtzman from Holtzman *et al.* 1997) with a CMD in which an additional component that consists of 8% of the original stars was placed uniformly between 10 and 15 kpc closer than the LMC along the line of sight. The two CMDs are nearly indistinguishable (this degeneracy in part motivated the suggestion of an intervening population near the LMC by Zhao (1998a)). The additional freedom regarding relative metallicities and reddenings for the foreground population would enable one to create even less distinguishable CMDs (Geha, priv. comm.). Therefore, although deep CMDs may eventually provide valuable constraints on the line of sight stellar distribution they are currently not able to confirm or refute the presence of stars within 15 kpc of the LMC at the fractional level proposed by ZL.

3.1.2. Gallart 1998

The multi-million star CMDs produced by the MCPS and the microlensing surveys provide a wealth of information on stellar evolution by containing statistically significant populations in rare phases of stellar evolution. An example of this claim is an overdensity that was identified by ZL along the giant branch (in BV CMDs with the

observational uncertainties of the MCPS, the red giant and asymptotic giant branches are indistinguishable). ZL associated this particular feature with the red giant branch bump (RGBB) seen in some globular clusters (King *et al.* 1985; Fusi Pecci *et al.* 1990). Gallart examined this feature in detail and identified it as the AGB bump that is predicted by the Padova isochrones (Bertelli *et al.* 1994 and references therein). She further demonstrated that the RGBB would appear at fainter magnitudes than the red clump in the LMC population. This revision of the identification of the feature along the red giant branch is important in terms of stellar evolution and star formation models of the LMC, but does not impact the hypothesis of an intervening population — except as a cautionary tale of the possible manifestation of previously unidentified stellar evolutionary phases.

Gallart also noted on the basis of synthetic CMDs that one should expect a population of stars extending vertically from the red clump for stellar populations with a component that is younger than 1 Gyr and that such a feature is observed in the Fornax and Sextans A dwarfs (Stetson 1997; Dohm-Palmer *et al.* 1997). This suggestion has important ramifications for the interpretation of the VRC, but we postpone exploring this option until we discuss the work by Beaulieu and Sackett (1998) which examined this possibility quantitatively.

3.1.3. Bennett 1998

A key step in determining if a candidate foreground population is an important source of uncertainty in the interpretation of τ_μ is the evaluation of the corresponding microlensing optical depth of such a population, τ_{fg} . For example, if the population suggested to be associated with the VRC has $\tau_{fg} \ll \tau_\mu$, then this result is a robust argument against the importance of this population because it is independent of the nature of the VRC. The origin of the VRC, at least with respect to microlensing surveys, becomes academic.

The conversion of a VRC surface density into a corresponding τ_{fg} is uncertain because the VRC is only a tracer of the possible foreground population. To convert the number of VRC stars observed into a surface mass density of the associated stellar population requires assumptions about the completeness of the survey, the distribution of distances among the stellar populations, and, most importantly, the mass-to-RC star ratio of the population (which in turn depends on the initial mass function (IMF) and the star formation history).

ZL adopt an empirical approach and assume that the proposed foreground stellar population is the same as the LMC population (eg. same M/L). They calculate the mass density per red clump star for the LMC population using the LMC rotation curve to derive the mass. Concerns have been raised about this approach (see the discussion of Gould (1998) below). Once the mass density per red clump star

for the LMC is known, ZL evaluate the surface mass density of the foreground population by attributing the same mass density per red clump star to each VRC star (additional geometric considerations were also applied). Once the surface mass density of the foreground population is evaluated the conversion to an optical depth is noncontroversial. ZL conclude that within large uncertainties (at least 50%) the observed lensing could be due to the foreground population, thereby potentially making this stellar population a critical component of the entire lensing budget along the line of sight to the LMC. Because of various “factor-of-two” uncertainties, this calculation does not prove that the proposed foreground population accounts for the observed lensing, but instead provides a viable plausibility argument. Given our philosophy (*cf.* §1.1), this result is sufficient for us to question the existence of halo MACHOs.

Bennett critiqued the calculation of τ_{fg} and presented an alternate analytic method for calculating the mass density of the foreground population. For a given IMF and star formation history, one can calculate the number of RC stars and the mass density of the entire population. Bennett presented such a calculation and concluded that *even if* the VRC were truly a tracer of a foreground population, such a population would *at most* contribute $\tau_{fg} = 0.13\tau_{\mu}$. This result directly contradicts the ZL calculation by at least a factor of several, perhaps a factor of 10, and argues that the VRC, regardless of its origin, is at most a small factor in the interpretation of the results from the microlensing surveys.

Because Bennett’s conclusion is critical in assessing the nature of the lensing sources, we examine the calculation in detail. His conclusion is robust *if* it is independent of assumed parameters, within the plausible range of such parameters. For example, if a slight change in the adopted IMF slope results in $\tau_{fg} \sim \tau_{\mu}$, then this approach is indeterminate and does not provide proof against the possible importance of a foreground population traced by the VRC.

Because this theoretical approach requires one to know the absolute number of VRC stars, rather than the ratio between VRC and LMC red clump stars as used by ZL, we need an accurate census of VRC stars. Bennett adopted the number of $\sim 70,000$ red clump stars in the ZL area ($2^{\circ} \times 1.5^{\circ}$) and a relative ratio between VRC and RC stars of 5% to derive a surface density of 1200 “foreground” stars per sq. degree. These are rough estimates from the ZL histograms and here we take a more precise census. There are 89849 RC stars for an RC defined to have $0.7 \leq B - V < 1.15$ and $18.85 < V < 19.5$. The area covered is 2.93 sq. degrees. The VRC fraction was estimated by ZL to be between 5 and 7%, but that number is sensitive to the model that is used to subtract the red clump contribution from the VRC and to the definition of the color and magnitude boundaries for the VRC stars. As can be seen in ZL’s Figure 4, the red clump model subtracts a substantial fraction of the potential VRC stars. Nevertheless, using 0.07 for the VRC/RC ratio, which was ZL’s upper limit (Bennett used 0.05 and Beaulieu and Sackett estimate it to be 0.08), we obtain a surface density

of 2,150 “foreground” stars per sq. degree. If instead, we use the definition of the VRC that is used elsewhere in this paper ($2.85 < C < 3.57$, $17.8 < V < 18.5$, where $C = 0.565(B - I) + 0.825(U - V + 1.15)$, *cf.* ZL), then we measure a surface density of 2,471 “foreground” stars per sq. degree (and derive $\text{VRC}/\text{RC} = 0.08$). These surface densities are 80% to 106% larger than the number adopted by Bennett and simply illustrate one of the many “factor of two” uncertainties that affect this argument, even before uncertainties in IMF’s and star formation histories are considered.

With the measurement of the number of red clump stars in hand, we can calculate the corresponding total number and mass of stars in this population. That mass can then be converted into a microlensing optical depth. Despite some problematic assumptions (such as the characterization of the LMC population as a single age population), we present the equivalent derivation to Bennett’s calculation, with the exception that we leave the IMF slope as a free parameter. From Bennett, the total mass in stars in a given population that is characterized by a single current main sequence turnoff mass given by $m_{to}M_{\odot}$ is given by

$$M = A \left(\int_0^{m_1} m_1^{\Gamma} dm + \int_{m_1}^{m_{to}} m^{\Gamma} dm + \int_{m_{to}}^{m_2} m^{\Gamma-1} m_{wd} dm \right), \quad (1)$$

where A is an arbitrary normalization constant, m_1 is the mass at which the IMF makes the transition from a power-law IMF to a flat IMF (*e.g.*, for $m < m_1$, $n(m)$ is constant) in solar masses, m_2 is the upper mass cutoff for the IMF in solar masses, m_{wd} is the mass of objects that have evolved past the horizontal branch and is taken to be $m_{wd} = (0.15m + 0.38)M_{\odot}$ (Iben and Renzini 1983), and we have expressed the IMF power-law slope as Γ . As defined by Scalo (1986) and adopted by Massey *et al.* (1995), $\Gamma = d \log \xi(\log m) / d \log m$, where $\xi(\log m)$ is the number of stars born per unit logarithmic (base ten) mass interval per unit area per unit time. For reference, a Salpeter (1955) mass function has $\Gamma = -1.35$. Equation (1) can be expressed as

$$M = A \left(\frac{\Gamma}{\Gamma + 1} m_1^{\Gamma+1} + \frac{0.85}{\Gamma + 1} m_{to}^{\Gamma+1} - \frac{0.38}{\Gamma} m_{to}^{\Gamma} + \frac{1.5\Gamma + 0.38(\Gamma + 1)}{\Gamma(\Gamma + 1)} 10^{\Gamma} \right). \quad (2)$$

The number of horizontal branch (red clump) stars, N_{RC} , is given by the mass range of stars that are currently sufficiently old to be on the horizontal branch. Bennett parameterizes that mass interval as ranging from m to $m(1 + \delta)$ where $\delta = t_{RC} / (3.75 t_{ms})$. For an adopted parameterization of the main sequence lifetime, $t_{ms} \propto m^{-3.75}$, and a horizontal branch lifetime, t_{RC} ,

$$\begin{aligned} N_{RC} &= A \int_{m_{to}}^{(1+\delta)m_{to}} m^{\Gamma-1} dm \\ &= A m_{to}^{\Gamma} \delta. \end{aligned} \quad (3)$$

Assuming a constant t_{RC} of 10^8 yrs (following Bennett and adopted from Castellani, Chieffi, & Pulone 1991) and $t_{ms} = 1.1 \times 10^{10} m_{to}^{-3.75}$ yr, then

$$N_{RC} = 0.00242 A m_{to}^{\Gamma+3.75}. \quad (4)$$

Taking Equation (1) and dividing by Equation (4) to obtain the total stellar mass per red clump star gives ,

$$\begin{aligned} M/N_{RC} = & \frac{413\Gamma}{\Gamma+1} m_1^{\Gamma+1} / m_{to}^{\Gamma+3.75} + \frac{351}{\Gamma+1} m_{to}^{-2.75} - \frac{157}{\Gamma} m_{to}^{-3.75} \\ & + \frac{620\Gamma + 157(\Gamma+1)}{\Gamma(\Gamma+1)} 10^\Gamma m_{to}^{-\Gamma-3.75}. \end{aligned} \quad (5)$$

The optical depth is calculated by multiplying the surface mass density in units of solar masses per square degree by $3.95 \times 10^{-14} (1-x)/x$ where x is the fractional distance of the lensing population relative to the source population, for an assumed distance to the LMC of 50 kpc. Therefore, the total optical depth is given by multiplying the surface number density of VRC stars, which in the foreground population hypothesis are the RC stars of this population, by M/N_{RC} and by $3.95 \times 10^{-14} (1-x)/x$.

To obtain numerical estimates for the optical depth we need to adopt numerical values for m_1 , m_{to} , Γ , and x . Bennett examined two values of m_1 , 0.3 and 0.6, to represent a Galactic globular cluster type population and a Galactic disk type population, respectively. The IMF slope Γ was chosen to be -1.3 , where $n(m) \propto m^{\Gamma-1}$ for $m_1 < m < 10M_\odot$. Selecting these values, and $x = 0.7$ corresponding to a distance for the foreground population of 35 kpc, the optical depth is $< 4 \times 10^{-8}$ for $0.9 < m_{to} < 1.7$ in exact agreement with Bennett’s Figure 1, and is much less than the observed value of $2.9_{-0.9}^{+1.4} \times 10^{-7}$ (Alcock *et al.* 1997a) or the smaller, less certain, measurement of Ansari *et al.* (1996) of 8.2×10^{-8} . Analysis of the combined data concludes that $\tau_\mu = 2.1_{-0.8}^{+1.3} \times 10^{-7}$ (Bennett 1998).

We now probe the sensitivity of this result on the adopted parameters. First, we examine the dependence on m_1 . As m_1 decreases, τ_{fg} increases because there are more lower mass stars that act as lenses. For example, a choice of $m_1 = 0.1$ leads to an increase in the optical depth of about 70%. Examining the reference cited for the choice of m_1 (De Marchi & Paresce 1997), the proper choice of m_1 does not appear to be precisely determined. Depending on the adopted mass-luminosity relationship for low mass stars, the mass function flattens somewhere between 0.3 and $0.13M_\odot$.

Second, we probe the dependence of the calculated τ_{fg} on the choice of Γ . Highly negative values of Γ (steeper IMFs) increase the optical depth because there are more low mass stars for each detected high mass star. Bennett adopted the standard value and one that appears to be consistent with observations of LMC associations ($\Gamma = -1.3 \pm 0.3$; Massey *et al.* 1995). However, the field IMF in the LMC appears to be significantly steeper, with some estimates going as far as $\Gamma = -4.1 \pm 0.2$ (Massey *et al.* 1995). If the appropriate slope is that steep, we in fact *overestimate* the observed

optical depth by a factor of 2 (even with $m_1 = 0.3$ and Bennett’s lower number for the VRC surface number density). More moderate values of the IMF slope, such as $\Gamma = -2$ (with, for example, $m_1 = 0.2$ and the somewhat higher number density of VRC stars we measure) give optical depth estimates in agreement with those of ZL ($\sim 0.5\tau_\mu$), and therefore suggest that the foreground population *could* be an important contributor to the total optical depth.

Admittedly, we have maneuvered the parameters in favor of high optical depth and the observational determinations of the field IMF in the LMC are valid only for high mass stars, but the selected parameters are well within the ranges currently allowed by observations. Even if $\Gamma = -1.6$ (a 1σ excursion from the Massey *et al.* ’s result for LMC OB associations) $m_1 = 0.2$, and $m_{to} = 1M_\odot$, $\tau_{fg} = 1 \times 10^{-7}$, which is roughly between a third and half of the observed optical depth (Alcock *et al.* 1997a; Bennett 1998). A combination of a comparable population behind the LMC (cf. Zhao 1998b), a warped Galactic disk (Evans *et al.* 1998), and increase contribution from the known lensing components (*e.g.*, LMC halo and flat galactic disk; Aubourg *et al.* 1999) *could* account for the difference between this optical depth and τ_μ .

For allowed parameter ranges, τ_{fg} can range from insignificant to $\sim \tau_\mu$. We find that the uncertainties in the IMF slope and the stellar mass at which the IMF may flatten, preclude the use of this argument to categorically conclude that a foreground population that has a contrast of $\sim 5\%$ in red clump stars cannot account for about half of the observed optical depth. We stress that Bennett’s adopted parameters and low optical depths are plausible, perhaps even preferable, but not unique.

As a final note, it is illuminating to calculate the lowest fractional population of foreground red clump stars that can provide at least half of the observed optical depth for various limiting parameter cases (*e.g.*, how precisely do we need to exclude a foreground population to exclude it as a major source of microlenses). For example, with $\Gamma = -4$, and the other parameters unchanged from those adopted by Bennett, the foreground population needs only to be 0.9% of the red clump stars. Therefore, even if 87% of the VRC stars are due to another phenomenon (*e.g.*, stellar evolution), there are still sufficient foreground stars to account for half of the observed optical depth. For $\Gamma = -2$, $m_1 = 0.1$, and $m_{to} = 1$, only 45% of the VRC stars need be foreground stars for $\tau_{fg} = 0.5\tau_\mu$. It will be exceedingly difficult to exclude such a small fraction of foreground stars without direct distance measurements to the lenses. The more extreme values of Γ are almost certainly inappropriate over a large mass range (for example, they would predict very large M/L), but the appropriate value is not well determined.

3.1.4. Gould 1998

Gould examined whether ZL’s implied M/L ’s, both for the LMC disk and the foreground population are reasonable and whether there is any existing evidence for tidal streamers/tails superposed on the LMC. The only existing photometric survey that is relatively deep over a sufficiently large area is the photographic survey of de Vaucouleurs (1957). Based on the surface brightnesses measured in that survey and the surface mass density necessary to reproduce the observed lensing optical depth, Gould concluded that the implied M/L of 12 was unacceptably high (if one was to account for the lensing by appealing to a stellar population that resembles known stellar populations). Furthermore, he concluded that de Vaucouleurs’s maps shows no evidence of a component (with sufficient surface mass density) extending beyond the LMC and concluded that either (1) there is no such component, (2) it is concentrated on the LMC and has a surface brightness profile similar to the LMCs, or (3) it is uniform over the entire area of the survey and so, much more extended than the LMC.

In addition to the standard difficulties in low surface brightness measurements with photographic material, the uneven foreground emission is, according to de Vaucouleurs (1957), “particularly serious”. Because of “galactic structures ... and the general luminosity gradient toward lower latitudes”, de Vaucouleurs was forced to apply a set of corrections that were constructed to remove irregularities “as well as possible by means of field corrections” and also to apply “considerable smoothing ... to reduce spurious details.” Because of such difficulties, de Vaucouleurs’s concluded that his procedure “effectively excludes the very weak outer extensions”. Given this irreversible treatment of the data, it is difficult to ascertain the precision of the outer surface brightness contours and the reliability with which tidal features may be identified (de Vaucouleurs used those data to determine a total magnitude and a surface brightness profile for the LMC, so his analysis did not require precise outer isophotes). Indeed, current studies are beginning to find signatures of possible stellar tidal structures near the LMC (Majewski *et al.* 1998; Geisler *et al.* 1998).

Gould compared the M/L derived using the mass density from ZL and the luminosity from de Vaucouleurs’s V -band measurement, converted to R for an assumed $V - R = 0.5$, to the I -band M/L given by Binney and Tremaine (1987) for the local Galactic disk (1.8 in solar units). We can redo that calculation using our photometry to avoid the issues related to the photographic data and the assumed color conversions. Our calculation has the added advantage that it provides the luminosity at the position in the LMC being discussed. We merge our data with the deep HST luminosity function (available for V and I from Holtzman’s data) and measure the V -band luminosity of the ZL region to be $5.1 \times 10^7 L_\odot$ (for a distance modulus of 18.5 and $M_{V,\odot} = 4.8$). This luminosity is a slight underestimate because the HST data are incomplete at faint magnitudes ($M_V > 7$), but those stars contribute little to the total luminosity. The corresponding average surface brightness of the ZL region is 22.9

$L_{\odot}\text{pc}^{-2}$. The average reddening, $E(B - V)$, measured for O and B stars in this area of the LMC is 0.2 mag (Harris, Zaritsky, & Thompson 1997) and includes Galactic reddening. For a standard interstellar extinction law (Schild 1977), which is valid at optical wavelengths for dust in the LMC (Fitzpatrick 1985), $A_V = 3.2E(B - V)$, so $E(B - V) = 0.20$ mag corresponds to $A_V = 0.64$ mag. After making this correction, the extinction-free V-band surface luminosity is $41L_{\odot}\text{pc}^{-2}$. This value is a slight overestimate if the VRC stars are indeed in the foreground (because their dereddened luminosity was included in the measurement), but the foreground population is $< 10\%$ of the LMC population so the effect is small. The ZL mass estimate, which corresponds to $103 M_{\odot} \text{pc}^{-2}$, implies $(M/L)_V = 2.5$.

We can also do the same calculation for the I -band, which is less affected by dust. The I -band luminosity of the ZL region (for stars with $I \leq 19.0$) is $3.2 \times 10^7 L_{\odot}$. Again using Holtzman’s data to correct for missing fainter stars and an I -band extinction correction of 0.38 mag, the surface brightness is $32 L_{\odot}\text{pc}^{-2}$ and $(M/L)_I = 3.2$. Finally, we note that the ZL region is not as luminous as other regions (*e.g.*, the left side of Figure 1) and so the global M/L for the LMC may be lower.

We compare these values of M/L to measurements of M/L for other galaxies. The calculated $(M/L)_V$ is not beyond the range observed for spirals (cf. Kent 1986; although his values are for the r -band and some slight color correction is necessary). The I -band values are easier to judge because many measurements exist for either the local disk ($(M/L)_I = 1.8$ as adopted by Gould from Binney & Tremaine) or for large samples of galaxies. Vogt (1995) presents $(M/L)_I$ for a wide range of spiral galaxies. Although Hubble types Sd and Sm are not well-represented in her sample, we adopt the values for Sc types as appropriate (this is possible because no strong type or luminosity dependence is found). The median $(M/L)_I$ (converted to $H_0 = 75 \text{ km sec}^{-1} \text{ Mpc}^{-1}$) is 2.5 and the upper quartile begins at $(M/L)_I = 3.1$. The median value for Sb galaxies (appropriate for our galaxy) is 2.0, in excellent agreement with the value of 1.8 from Binney & Tremaine. In comparison to these results, $(M/L)_I$ derived for the LMC is slightly larger than expected, but not beyond the range of values observed for other galaxies (it lies just inside the upper 25%). Given the reasonable agreement in the inferred LMC M/L ratios for both V and I and determinations for other galaxies, and the uncertainties in the multiple steps required for this comparison, we find no reason to conclude that ZL’s inferred LMC mass, and by association that of the foreground component, can only be explained with a dark matter dominated (*e.g.*, $M/L > 12$) stellar population. Again, we stress that “factor of two” effects are ubiquitous and can work in either direction (for example, ZL adopted a disk inclination of 33° for the LMC but published values range from 22° (Kim *et al.* 1998) to 48° (Bothun & Thompson 1988) which result in velocity inclination corrections that vary by a factor of two). We conclude that there are large uncertainties in the derived masses, but that masses as large as implied by ZL’s analysis are not sufficiently extreme to be

confidently excluded.

Finally, Gould noted that ZL exclude the contribution of the dark matter in the LMC to the rotation curve and thereby inflate the stellar M/L (and so overestimate the lensing contribution of a standard stellar population). Kim *et al.* (1998) recently published and analyzed a high resolution H I map of the LMC. They present mass models and conclude that no dark matter halo is required to fit the H I data. To fit their H I observations *and* the carbon star data at larger radii (Kunkel *et al.* 1997), they do use a model that includes a dark matter halo. From their Figure 6, we infer that the halo’s contribution to the mass inside 3 kpc is $< 20\%$. Therefore, although while Gould’s assertion is strictly true, the discrepancy is apparently minor given the “factor-of-two” uncertainties present elsewhere. For their global model, Kim *et al.* calculate that the stellar $(M/L)_R$ is 1.8 (where the luminosity is taken from de Vaucouleurs (1957)), which is less than a factor of two discrepant with either the V or I -band M/L ’s derived above. Without a dark matter halo (or with a smaller contribution from the dark matter halo within 4 kpc), the stellar M/L would increase by up to $\sim 20\%$ and be in even better agreement with our estimates of M/L .

3.1.5. Johnston 1998

ZL suggested that one possible source of foreground stars was the suggested tidal streamer (Zhao 1998a) that arose as a result of the LMC-Milky Way interaction in a previous perigalacticon passage. Johnston examined this suggestion in greater detail and concluded that either such a feature would lead to unacceptably high faint star counts across a large section of the sky or that the feature would disperse quickly ($\sim 10^8$ years). Her study does not exclude other possible sources of foreground stars such as tidal material from an SMC-LMC interaction or from a denser than expected LMC halo. The SMC-LMC interaction that is thought to have occurred within the last $\text{few} \times 10^8$ years (Gardiner & Noguchi 1996) may be an attractive candidate for the source of tidal material along the line of sight. The environment around the LMC-SMC is sufficiently complicated (*cf.* Putnam *et al.* 1998) that the presence of stars outside the body of the LMC with a total mass of a few percent of the LMC should not be surprising.

3.1.6. Beaulieu and Sackett 1998

The study by Beaulieu and Sackett provides the strongest challenge yet to the interpretation of the VRC as a foreground population by suggesting that the young (< 1 Gyr) LMC stars evolve into a superluminous red clump phase that populates the VRC region of the CMD. This argument is bolstered by theoretical isochrones for such stars, the clear presence of young main sequence stars in the LMC, the identification of such superluminous red clump stars in the Galactic HIPPARCOS data, and the

identification of such stars in the Fornax and Sextans A dwarf galaxies (Stetson 1997; Dohm-Palmer *et al.* 1997). Quantitatively, they support their suggestion by presenting models for the star formation history of the LMC that produce the observed number of stars in the VRC region of the CMD. However, recall from §3.1.3 that to the limit of our current understanding of the IMF, it is possible that only a minor fraction ($< 50\%$) of the VRC stars need be foreground stars to account for a large fraction of τ_μ . Therefore, a precise estimate of the fraction of VRC stars that are young red clump stars is necessary.

If the star formation history adopted by Beaulieu and Sackett is correct, then their models predict that any foreground population contributes at most only slightly ($< 25\%$) to the VRC. However, once again the models depend on assumption regarding complex issues such as the star formation history and IMF. We examine the dependence of those models on some of the selected parameters.

We adopt the Bertelli *et al.* (1994) isochrones, a standard (Salpeter) IMF slope, and add observational uncertainties to generate CMDs corresponding to a variety of star formation histories. From these data we have recreated the histogram of differential magnitude relative to the red clump centroid for the narrow color cut around the RC that was presented by ZL. We have selected three star formation histories for comparison: (1) a constant star formation rate (SFR) over the most recent 1 Gyr, and nothing prior, (2) a constant SFR over the most recent 1.3 Gyr and nothing prior, and (3) the star formation history presented by Geha *et al.* (1997) based on analysis of deep HST CMDs.

There are two representative tests of the models. First, the ratio of the number of main sequence stars with main sequence lifetimes ~ 0.8 Gyr to the number of VRC stars constrains whether the model is able to generate VRC stars provided that there has been a recent episode of star formation. This ratio was used by Beaulieu and Sackett to demonstrate the feasibility of their hypothesis. Second, the ratio of VRC to RC stars tests whether the entire star formation history is consistent with the observed number of VRC stars relative to RC stars. If the first test is not satisfied, then one can conclude that this phase of stellar evolution (young, massive RC stars) cannot populate the VRC in sufficient numbers, regardless of the star formation history. If the first test is satisfied, but the second is not, then one concludes that this phase of stellar evolution can populate the VRC in sufficient numbers, but that the recent star formation rate was insufficient to account for the observed number of VRC stars. Beaulieu and Sackett demonstrated that the first test is satisfied (their models were able to reproduce between 75% and 100% of the ratio of main sequence stars to VRC stars).

We present the histograms from vertical (constant color) cuts through the simulated CMDs at the position of the red clump in Figure 6. Our first conclusion is that stellar evolution is able, in principle, to populate the VRC. Our second conclusion is that

a model with a constant star formation over the last 1.3 Gyr (but no previous star formation) is at least one model that reproduces the observed ratio of VRC to RC stars. This conclusion confirms the result of Beaulieu and Sackett that a model with a constant SFR over the last $\sim$ Gyr is able to generate the appropriate number of VRC stars. Our third conclusion is that a model with an LMC star formation history independently derived from HST CMDs will significantly *underproduce* the number of VRC stars (defined to have $-1.4 < \Delta m_V < -0.7$) relative to the RC stars (defined to have $-0.325 < \Delta m_V < 0.325$) within this color cut (for the Geha *et al.* star formation history, the calculated ratio is only 26% of the observed ratio). We conclude that given the uncertainties in the star formation history, stellar models (*e.g.*, overshooting which affects the lifetime), and binarity (which we did not explore, but which may provide additional free parameters), we are unable to unequivocally determine whether stellar evolution does or does not populate the VRC region of CMDs in sufficient numbers, but that a model with a “realistic” star formation history defined independently of the VRC falls short by about a factor of two. Our understanding of the star formation history of the LMC is inadequate to be able to reach definitive conclusions about the relative fraction of young red clump stars that constitute the VRC with better than a 50% uncertainty. We will attempt to constrain this hypothesis with our new data in §4.

3.2. Summary of Previous Studies

We close this section by stressing that while certain “reasonable” assumptions can lead one to conclude that foreground stars are at most a small fraction of the VRC stars or that even if most of the VRC stars are foreground RC stars this population will not contribute significantly to τ_μ , these conclusions depend sensitively on adopted assumptions. *A foreground population that is responsible for a significant fraction of τ_μ is consistent with the available data.* If it is so difficult to exclude a relatively well-defined, spatially extended stellar population, it will be nearly impossible to exclude more subtle populations such as a slightly thicker LMC (Graff *et al.* 1998) or a population less than 10 kpc behind the LMC (Zhao 1998b) without having direct distance measurements to the lenses.

4. New Constraints on the Nature of the VRC

We set out to further constrain the nature of the VRC stars in two ways: (1) by obtaining stellar photometry over a larger area of the LMC to examine the distribution of VRC stars relative to other LMC stellar populations, and (2) by measuring VRC and RC radial velocities to place kinematic constraints on the population. The data acquisition was discussed in §2, we now discuss the results from each of those datasets below.

4.1. Spatial Distribution

The spatial distribution of the VRC stars can provide a fundamental clue to the nature of the VRC stars. If the VRC stars are seen exclusively in sites of recent ($\lesssim 1$ Gyr) star formation, then the Beaulieu and Sackett model would be confirmed. Alternatively, if VRC stars are found in regions where there are no comparably aged stars, then VRC stars are not exclusively young red clump stars. Therefore, a simple test is to compare the spatial distribution of VRC stars and main sequence stars that are of approximately the same mass as the VRC progenitors.

The VRC stars here are defined in color as in ZL and in magnitude as by Beaulieu & Sackett for more direct comparison (the same color criteria as Beaulieu & Sackett applied could not be used because we do not have R -band data). We define the VRC stars to have $2.85 < C < 3.57$ and $17.8 \leq V \leq 18.5$. We define the main sequence (MS) stars to which we compare to have $-0.2 < B - V < 0.3$ and $19.15 < V < 19.5$ (the same magnitude range as chosen by Beaulieu & Sackett). The different color criteria between our definitions and those of Beaulieu & Sackett have a minimal effect. For example, their preferred model results in a ratio of VRC stars to MS stars of 0.23 for their selection parameters and our comparable model (constant star formation for the last 1.3 Gyr and no star formation prior to that) results in a ratio of 0.20 for our selection parameters. We bin our data into $6' \times 6'$ squares (comparable to the area of a single image in the Beaulieu and Sackett study).

In Figure 7 we compare the stellar density distributions of VRC and MS stars. For the assumption adopted by Beaulieu & Sackett of constant star formation rate over the brief time separating the formation of the VRC progenitor stars and the current MS stars, the ratio of VRC to MS stars should be a constant across the area. From the Figure it is evident that there is some correlation between the two (*e.g.*, the enhancement of VRC stars in the lower left and upper right of the area), but the VRC distribution appears more uniform than the MS distribution. For a quantitative comparison we divide the VRC image by the MS image and examine this ratio on a pixel-by-pixel basis. The distribution of those ratios as a function of MS stellar density is plotted in Figure 8.

The systematic variation of the mean ratio with MS star density indicates that the number of VRC stars is not directly proportional to the number of MS stars. Variations of this ratio could be due to a variation of the star formation rate as a function of location over the time in which the current VRC stars were formed and the current MS stars were formed, to the greater diffusion of the slightly older VRC stars from their star formation sites, or to the presence of an additional component in the VRC that is uncorrelated with the MS stars. Nevertheless, variations seen in Figures 7 and 8 demonstrate (1) that the measured ratio in any single $6' \times 6'$ box cannot be directly compared to the ratio calculated from simulated CMDs and (2) that at least some VRC stars are associated with recent star formation regions.

Regardless of the correct explanation for the variation in VRC/MS, this ratio must approach that estimated from the synthetic CMDs when measured over a large region of the LMC if the young RC hypothesis is correct. The average of VRC/MS over our entire field is 0.31, while the models (using either the model with constant star formation over the last 1.3 Gyr or the Geha *et al.* star formation history) predict 0.20 and 0.14, respectively. The comparison to the “realistic” star formation history model suggests that over the entire area $\sim 50\%$ of the VRC stars are not related to young red clump stars and that the behavior of VRC/MS is not entirely attributable to either of the first two possibilities.

The existence of a uniform VRC component can be quantified using a simple model for the ratio of VRC stars to MS stars. If the young red clump component of the VRC stars is an unknown fraction, f , of the MS stars and the remainder is in a uniform component of A stars per bin, then

$$\frac{VRC}{MS} = \frac{A + f \cdot MS}{MS}. \quad (6)$$

Fitting this equation to the data presented in Figure 8, we derive $f = 0.088$ and $A = 7.25$, with $\chi^2 = 1.2$. This result implies that 28% of the VRC stars are young red clump stars and that the remainder is in a uniform component, which *could* be a foreground population. Alternatively, one can take f as given from stellar population models. For example, from the Geha *et al.* star formation history, $f = 0.14$. Fitting such a model we derive $A = 4.88$ with $\chi^2 = 2.9$. While the latter is a poorer fit (which can be excluded with 90% confidence), it still predicts that only 45% of the VRC stars are young red clump stars. Models that predict that a larger fraction of VRC stars are young red clump stars can be discriminated with higher confidence.

As discussed above, the possibility that VRC and MS stars have different clustering scales can affect this analysis. Variable clustering scale may be present if the young red clump stars have had sufficient time to diffuse farther away from sites of star formation on average than have the MS stars. To check if this is a substantial contributor to the previous result, we redo the analysis using boxes that are $3' \times 3'$ and $12' \times 12'$. The results are shown in Figure 9 and are consistent with the previous best fit relation. The best fit model to the data from the $3' \times 3'$ boxes suggests that 21% of the VRC stars are associated with the MS stars (instead of 28% as derived from the best fit to the data from the $6' \times 6'$ boxes). This difference is in the sense expected if the VRC and MS stars are distributed similarly but the VRC stars have slightly larger clustering scales. Nevertheless, this difference does not affect the conclusion that a large fraction of the VRC stars are unassociated with the MS stars. The distribution of VRC/MS from boxes ranging from $3'$ to $12'$ on a side give consistent results that a minority fraction ($< 50\%$) of the VRC stars are directly correlated with the main sequence stars. Although the simplest interpretation of the spatial modeling is consistent with the results from the simulated CMDs, other interpretations involving variable star

formation or more complex dynamical evolution are possible.

Why does our conclusion regarding the origin of the VRC differ from that of Beaulieu & Sackett? Over a wide range of MS star densities the ratio of VRC/MS in $6' \times 6'$ areas is ~ 0.2 . It is therefore not unexpected that when Beaulieu and Sackett analyzed four fields of this size, three of those had ratios in that range. On average (weighing their four fields equally), their measurement of VRC/MS is 0.33, which is nearly identical to our “global” average of 0.31. Therefore, the different conclusions are not due to differences in the definition of the VRC or MS stars (or to gross differences arising from reddening or completeness corrections), but rather in our ability to explore a wide range of MS star densities, average over many fields to improve our statistics, and on our use of the Geha *et al.* star formation history. Clearly, the range of possible star formation histories is infinite and it is quite possible that there are many plausible histories that produce a sufficient number of young red clump stars, but the detection of dark matter should not rest on such ill-constrained models.

4.2. Radial Velocities

The radial velocity distribution of VRC stars is shown in Figure 10. The small number of stars at ~ -275 km sec $^{-1}$ are Galactic stars and they illustrate that even though the location of the VRC in the CMD places it near the locus of Galactic contamination (disk stars), the level of contamination is minor. The bulk of the VRC stars are centered at 0 km sec $^{-1}$ rather than 275 km sec $^{-1}$, which is an artifact of how the velocity template was generated. The velocities for the red clump stars are presented in Figure 11, and the similar centroid velocities for the VRC and RC demonstrate that the VRC stars have an average velocity that is statistically indistinguishable from that of the LMC. This result confirms the result of Ibata, Lewis, & Beaulieu (1998) based on 16 VRC stars.

As Ibata *et al.* argue, the agreement between the VRC and LMC velocities argues for a close relationship between the VRC and LMC stars and almost certainly excludes the possibility that the VRC is due to a foreground population unrelated to the LMC (such as an intervening galaxy or tidal streamer). Therefore, any interpretation of the the VRC as foreground material, is now constrained to place these stars in close association with the LMC (they must be either gravitationally bound or recently tidally removed).

We proceed to measure the velocity dispersion of the VRC stars in two ways. The first approach we use is an iterative maximum likelihood fitting algorithm that fits to the unbinned distribution of data. We have eliminated stars with $|\Delta v| > 100$ km sec $^{-1}$ and with $R < 6$. These cuts eliminate galactic contamination and unreliable velocity measurements. The results from the remaining 190 VRC stars are $\bar{v} = 1.2 \pm 3.2$ km sec $^{-1}$ and $\sigma = 18.4 \pm 2.8$ km sec $^{-1}$ (95% confidence interval used throughout for

uncertainties in mean velocities and velocity dispersions). The second approach we use fits a Gaussian to the binned data. By convolving the underlying distribution, characterized by a mean velocity, $\bar{v}$, and dispersion, σ , with the distribution of observational errors, we generate the expected distribution of radial velocities. With a standard definition of χ^2 (the sum of the differences squared between the histogram and the convolved expectation divided by the uncertainties from Poisson statistics squared), we associate a χ^2 with each model and determine the best fitting model and allowed parameter ranges. For the same sample of stars as used for the unbinned parameter fitting, and a bin size of 10 km sec⁻¹ bins we find $\bar{v} = 3.2^{+1.8}_{-4.4}$ km sec⁻¹ and $\sigma = 17.5^{+3.8}_{-3.7}$ km sec⁻¹ ($\chi_2 = 1.1$). The two approaches provide a consistency check on the algorithms used to determine the underlying dispersion given the substantial observational uncertainties. However, we adopt the results from the first method because the results do not depend on an adopted bin size. As is evident from Figure 10, the Gaussian model provides an acceptable fit. Finally, we attempt to correct for the fact that the two fields, even though they are close to each other, could have a different mean velocity due to the projection of the LMC’s rotation curve on the plane of the sky. Rather than rely on a model, we make the most favorable assumption for lowering the velocity dispersion, which is to set the mean velocities of stars in the two VRC fields equal. The resulting velocity distribution has $\sigma = 18.0 \pm 2.8$ km sec⁻¹. We conclude that the measured VRC line-of-sight velocity dispersion is > 15 km sec⁻¹ with 95% confidence.

For the red clump stars (Figure 11), we have larger observational uncertainties and fewer data (75 stars), so the kinematic constraints are less precise. The unbinned fitting results in $\bar{v} = -4.6 \pm 9.0$ km sec⁻¹ and $\sigma = 32.2 \pm 7.6$ km sec⁻¹. The mean velocity of the fitted Gaussian to the binned data is -6^{+21}_{-12} km sec⁻¹ and the dispersion is 32^{+19}_{-16} km sec⁻¹. The mean velocity calculated directly from the data (no assumed model) is -4.6 ± 9.0 km sec⁻¹ (1σ), which is in exact agreement with the parameters from the unbinned fitting algorithm.

We use the velocity dispersion vs. age relation for stellar populations given by Villumsen (1985),

$$\sigma_z = \sigma_{z,0}(1 + t/\tau)^{0.31}, \quad (7)$$

where $\sigma_{z,0}$ is the velocity dispersion of the cold gas, t is the age of the population, and τ is a “diffusion” timescale to estimate a mean “kinematic” age for the VRC stars. This relation was derived for Galactic stars. While we assume that the general physical behavior is the same, we normalize the relation for LMC populations. Adopting an initially vertical velocity dispersion of 5.4 km sec⁻¹ (from the H I gas) and 19.1 km sec⁻¹ for a population of planetary nebulae of mean age 3.5 Gyr (Meatheringham *et al.* 1988), we derive $\tau = 6.0 \times 10^7$ yrs, in close agreement with the value derived by Wielen (1977) of 5×10^7 yrs for a growth rate of $(1 + t/\tau)^{1/2}$ (for Galactic stars). Using the Villumsen age-velocity dispersion, we calculate that the mean age of the

VRC population is 3.1×10^9 years and that a 95% confidence lower limit on the age is 1.8×10^9 years (this estimate is only valid if the VRC stars are a single, disk population). Similar value are obtained using the original Spitzer and Schwarzschild (1951,1953) formula ($\sigma \propto (1 + t/\tau)^{1/3}$) or Wielen’s parameterization.

Does the age-velocity relationship provide reasonable and reliable estimates of the velocity dispersion for young stellar populations? To test whether young stellar populations distributed through the LMC do have a low line-of-sight ($< 15 \text{ km sec}^{-1}$) velocity dispersion we reanalyzed the velocity data obtained by Feast, Thackeray, & Wesselink (1960,1961) on supergiants in the LMC. We adopt the same sample of stars that they use in their dynamical analysis. Because these stars, which are typically O through A supergiants, are distributed over the entire LMC, the velocities have to be corrected for the rotation of the LMC. We take the simplest model (a flat disk with a constant rotation curve at all radii) to avoid using additional parameters to deflate the velocity dispersion, and use the mean inclination and semimajor position angle found by Kim *et al.* (1998) from the study of the H I kinematics ($i = 33^\circ$, a semimajor position angle of -10°). We fit the model to derive the rotation velocity, 42 km sec^{-1} , and the systematic velocity, 271 km sec^{-1} . We then subtract this model from the data and find the velocity dispersion to be $19.6 \pm 4.4 \text{ km sec}^{-1}$. However, three of the most deviant points from the mean rotation curve are at radii $< 1 \text{ kpc}$ for which the rotation curve is steeply rising (Kim *et al.*) and where the influence of the bar is probably not negligible. Three other stars are also on the extreme tails of the distribution and appear to artificially inflate the dispersion. The observed distribution does not appear to be Gaussian and a model distribution with the calculated dispersion overpredicts the number of stars at intermediate velocity differences and underpredicts the number at small velocity differences (see Figure 12). Eliminating the three stars near the center of the LMC and the three most extreme outliers leaves us with a distribution that is well fit by a Gaussian of velocity dispersion $10.0 \pm 3.8 \text{ km sec}^{-1}$ (note that the VRC velocity distribution shows no sign of extended non-Gaussian wings and so its dispersion does not appear to be inflated by outliers). Removing only the two stars at $\Delta v = 50 \text{ km sec}^{-1}$ results in $\sigma = 12.5 \pm 3.6 \text{ km sec}^{-1}$, which is still inconsistent with the measured VRC dispersion at the greater than 95% confidence level, but roughly the value expected for a 0.8 Gyr old population. Finally, we stress that the rotation model is highly simplified and that deviations, such as a systematic change in the line of nodes, warping, and a non-flat rotation curve have been observed (Kim *et al.*) and will cause us to overestimate the dispersion of these stars. We conclude that there are stellar populations that are sufficiently young and dynamically cold ($\sigma \sim 10 \text{ km sec}^{-1}$) and that such small velocity dispersions can be measured using velocity measurements that have uncertainties that are $\sim 10 \text{ km sec}^{-1}$.

We can further attempt to constrain the VRC population by modeling the velocity distribution as a sum of two Gaussians, one with the velocity dispersion expected for

a population of age ~ 0.8 Gyr and the other undetermined. By fitting the range of relative amplitudes, we can place a limit on the fraction of the VRC population that could be in a cold, young disk component. We fix the velocity dispersion of the young component at 12.3 km sec^{-1} (derived from the Villumsen dispersion-age relation), and allow the means of the two Gaussian to vary. We find that models in which the young disk component contributes more than 77% of all the VRC stars can be excluded at $>95\%$ confidence (all models where the young population contributes $< 77\%$ are allowed, and for models with a high fraction of young stars the velocity dispersion of the undetermined component is $\sim 35 \pm 10 \text{ km sec}^{-1}$). While these results do not provide as strong a constraint as that provided by the spatial distribution, it is entirely consistent with the implications from the stellar population models and the spatial distribution that the young red clump stars contribute $\lesssim 40\%$ of the VRC stars.

The VRC stars have a lower velocity dispersion than the RC stars, and certainly lower than one would expect for a halo population of the LMC. Therefore, at first glance it appears that the VRC stars could not be a constituent of a dynamically hot component. However, we caution that because of the magnitude selection of VRC stars (*i.e.*, that they be brighter than the RC stars) we would have selected only those stars on the near side of the LMC “halo”. The measured velocities may not be a fair sampling of the velocity distribution of these stars and could produce a significant underestimate of the true velocity dispersion, if indeed they are distributed throughout the LMC halo.

There are at least two caveats with regard to this velocity analysis. First, velocity uncertainties are notoriously hard to estimate precisely. Because the measured velocity dispersion is not grossly larger than the velocity uncertainties, the resulting dispersion is *highly* sensitive to the adopted uncertainties. It is possible that the velocity dispersion is inflated by observational uncertainties. An increase in the velocity uncertainties by 50% lowers the best fit velocity dispersion to 12.8 km sec^{-1} and the 95% confidence limit on the velocity dispersion to 9.2 km sec^{-1} (at which point a young population could not be ruled out). Although we have no evidence that indicates that the velocity uncertainties are severely underestimated and our dispersion measurement is consistent with that of Ibata, Lewis, & Beaulieu (1998), the possibility exists. A second caveat is that giant branch stars may contaminate the VRC region. In particular, some contamination may be due to stars on evolutionary blueward loops (“blue noses”) near the AGB bump (Gallart 1998). Such stars are older than 1 Gyr and would have the necessary kinematics and uniform spatial distribution that are observed. Therefore, advocates of halo MACHOs can assert that the VRC consists of young red clump stars plus an older contaminating component, such as blue nose stars. We conclude that the composition of the VRC is mixed, but that a significant intervening population has not yet been ruled out.

4.3. Microlensing Revisited

In this discussion, we have suggested that some adjustments to the parameters adopted by ZL in their calculation of the optical depth might be warranted. For example, as many as half of the VRC stars may be young red clump stars rather than foreground stars. How does this affect the possible microlensing optical depth from such a population?

A straightforward calculation, along the lines of Gould (1998) goes as follows. In the ZL area there are 2471 VRC stars / sq. degree, the ratio of VRC to RC stars is 0.08 (as defined in §3.1.3), the ZL area is 2.93 sq. degrees, and the total I -band luminosity is $7.1 \times 10^7 L_\odot$ (corrected for extinction and stars fainter than the magnitude limit of the MCPS, *cf.* §3.1.4). We presume that half of the VRC stars are foreground stars. Assuming that the foreground stellar population is identical to the LMC stellar population, the luminosity of the foreground population is $2.8 \times 10^6 L_\odot$. For a typical $(M/L)_I$ for Sc galaxies (2.5; Vogt 1995), the total mass in this component over the ZL area of the LMC is then $7.1 \times 10^6 M_\odot$. The surface mass density is then $2.4 \times 10^6 M_\odot$ per sq. degree, or $5.0 M_\odot \text{ pc}^{-2}$ if the foreground stars are at 40 kpc. Using the same formula as used by Gould (rewritten for τ_{fg}),

$$\tau_{fg} = \frac{2.9 \times 10^{-7}}{47} \left(\frac{D}{10 \text{ kpc}} \right) \quad (8)$$

where $D \equiv d_{ol}d_{ls}/d_{os}$ and d_{ol} , d_{ls} , and d_{os} are the distances between the observer, the lensing structure, and the sources in the LMC, we calculate that $\tau_{fg} = 2.5 \times 10^{-8}$ ($\sim 9\%$ of τ_μ). This result agrees with Bennett’s (1998) and Gould’s (1998) conclusions that even if there is a foreground population it would not contribute significantly to τ_μ .

However, to demonstrate the effect of various uncertainties, we redo the calculation with slightly different (but plausible) parameters. First, we calculate the VRC/RC ratio using our entire dataset and obtain that the ratio is 0.096. Instead of adopting that 50% of the VRC is foreground, we will use the result from the best fit model described in §4.1 that 72% of the VRC is in a uniform component, which we will attribute to a foreground population. We will also use $(M/L)_I = 3.2$, which is consistent with what ZL derived and within the observed range of $(M/L)_I$ ’s for Sc galaxies (Vogt 1995), and a distance to the lenses of 35 kpc. For these parameters, we derive that $\tau_{fg} = 9.2 \times 10^{-8}$, which is 32% of τ_μ . If τ_μ is revised downward to $\sim 2 \times 10^{-7}$ (Bennett 1998), then this population can account for about 50% of the observed microlensing.

Finally, we stress that the foreground population does not need to produce $>50\%$ of τ_μ to be important. A foreground tidal or halo stellar population should have a counterpart behind the LMC. The lensing of that population by LMC stars will result in a comparable contribution to τ_μ (Zhao 1998b). In addition, if the Galactic disk contributes an optical depth as large as 1×10^{-7} (Evans *et al.* 1998), then the VRC

foreground population only needs to contribute $0.25\tau_\mu$ for these three populations to account for τ_μ . Finally, the “standard” intervening stellar components (as estimated by Alcock *et al.* 1997a) have $\tau = 0.5 \times 10^{-7}$, with significant uncertainties (*cf.* Aubourg *et al.* 1999) even prior to appealing to unknown tidal tails or tilted Galactic disks. Therefore, arguments against the importance of intervening populations on the interpretation of microlensing events should not consider a single population in isolation, but rather the sum of all plausible populations.

5. Summary

The microlensing surveys have convincingly demonstrated that microlensing can be detected. When the number of lensing events observed along the line of sight to the Large Magellanic Cloud (LMC) is compared to that expected in the standard model of the structure of the LMC and the Galaxy (Alcock *et al.* 1997a) there is an excess of lensing events. Several authors, including some of the authors of this paper, have suggested previously that the “excess” lenses may not be halo MACHOs but rather normal stars along the line of sight. In particular, Zaritsky & Lin argued that the color-magnitude diagram of the LMC has a vertical extension of the red clump (the VRC) that is consistent with the presence of a foreground population that lies within 15 kpc of the LMC. Various studies have attempted to refute almost every aspect of that claim.

We adopt a two step approach in revisiting this issue. First, we examine each of the rebuttal studies to determine whether the arguments presented can convincingly discriminate against the possible foreground population suggested by ZL. Second, we obtain new data to test the most damaging of the rebuttal studies (Beaulieu & Sackett 1998), which suggested that the VRC stars are all young, massive red clump stars.

From our examination of the published studies, we find that to within the observational and model uncertainties one cannot convincingly refute (or prove) the existence of a foreground population that exists within 0-15 kpc of the LMC along the line of sight and that provides a significant ($\sim 50\%$) fraction of the total microlensing optical depth. Therefore, we conclude that systematic uncertainties dominate the current efforts to measure the total mass in halo MACHOs. Additional lensing events, if they do not (1) dramatically alter the measured microlensing optical depth, (2) dramatically alter the spatial distribution of lenses across the LMC, or (3) provide a large number of anomalous lensing events for which transverse velocities can be calculated (Alfonso *et al.* 1998; Alcock *et al.* 1998, Rhie *et al.* 1999), will not allow us to reach a significantly different conclusion about the nature of the lenses. However, even if an intervening stellar population is responsible for a significant fraction of the microlensing events, some events may be due to halo MACHOS and these lenses could still be a fundamental component of the Galaxy. Direct distance estimates for lenses

will eventually avoid the complicated issues described here in understanding the line of sight stellar distribution.

From our new data and analysis, we find three reasons to question the suggestion that was quantitatively pursued by Beaulieu & Sackett (1998), that the VRC stars are exclusively young, massive red clump stars. First, synthetic color-magnitude diagrams created using a star formation history derived independently from deep HST data (Geha *et al.* 1998) suggest that $< 50\%$ of the vertical red clump (VRC) stars are young, massive red clump stars. Second, a detailed comparison of the VRC and young main sequence stars, suggests that $\lesssim 40\%$ of the VRC stars correlate spatially with the MS stars. The remainder are in a more uniform distribution, and so are unlikely to be young red clump stars. Lastly, our measurement of the velocity dispersion of 190 VRC stars ($18.4 \pm 2.8 \text{ km sec}^{-1}$ (95% confidence interval) is inconsistent with the expectation for a young ($\lesssim 0.8 \text{ Gyr}$) disk population (but consistent with that of the red giants). We confirm the measurement by Ibata *et al.* and agree with their conclusion, that even if the VRC stars are foreground stars they are unlikely to be from a stellar population unassociated with the LMC. We conclude that various arguments suggest that $\sim 40\%$ of the VRC stars are young, massive red clump stars. The remaining stars may be attributable to another stellar evolutionary phase or to an intervening population associated with the LMC (possibly an extended halo, stars that were recently tidally removed, or possibly a much thicker, more massive disk than previously expected (Aubourg *et al.* 1999)).

We apply a straightforward calculation for the microlensing optical depth that avoids some of the subtle issues of star formation histories and IMFs by relying only on an adopted M/L for the stellar populations, which we determine empirically from the studies of other galaxies. We find that for one set of parameters the optical depth from the possible foreground population (consisting of $\sim 50\%$ of the VRC stars) is $< 10\%$ of the optical depth measured by Alcock *et al.* (1997). For somewhat different parameters, we find that the microlensing optical depth from the allowed foreground population is $\sim 30\%$ of the Alcock *et al.* optical depth. More extreme, but not yet excluded, parameter choices can lead to percentages $> 50\%$. In combination with known stellar components and other possible sources of contamination (*e.g.*, a background population, a slightly thicker LMC, and more Galactic disk stars), this population can produce τ_μ . Whether it does, is an open question.

We have, given our stated motivation of strongly questioning whether the data yet demand the need for halo MACHOS, focused on attempting to remove published objections to the intervening stellar population model. Some intervening populations, such as an intervening stellar stream or dwarf galaxy have by now been excluded (cf. Alcock *et al.* 1997b and Ibata *et al.* 1998). We have argued here that a population associated with the LMC (either within the LMC halo or very recently tidally lost) has not yet been excluded. In the interest of some balance, we restate the basic arguments

against interpreting the VRC as an intervening population. First, stellar evolution models and plausible star formation histories are able to account for the number of VRC stars. Second, the measured kinematics of VRC stars do not show any grossly anomalous behavior (a difference in mean velocity with the LMC or non-Gaussian behavior) and therefore do not argue strongly against the VRC stars being in the LMC disk. Lastly, even if one accepts the interpretation of VRC stars as foreground (fg), plausible models for the conversion of the number of VRC stars into τ_{fg} result in $\tau_{fg} \ll \tau_{\mu}$. However, our goal was to demonstrate that as reasonable (and perhaps correct) as these arguments may be, numerous large uncertainties remain. Because of the importance of the nature of dark matter to a wide range of astrophysics, significant uncertainties must be removed before the case for its identification becomes compelling.

In conclusion, despite various investigations into the nature of the VRC population (including this one), we are only able to reiterate the closing sentence of ZL’s abstract — “We conclude that the standard assumption of a smoothly distributed halo population out to the LMC cannot be substantiated without at least a detailed understanding of several of the following: red clump stellar evolution, binary fractions, binary mass ratios, the spatial correlation of stars within the LMC, possible variations in the stellar populations of satellite galaxies, and differential reddening — all of these are highly complex.” To calculate the microlensing optical depth from such a population one must add to that list of problem issues the initial mass function and star formation history. Given these uncertainties, we find that the hypothesis of a foreground population within ~ 15 kpc of the LMC (which is intriguingly close to the LMC’s apparent tidal radius, $\gtrsim 13$ kpc; Schommer *et al.* 1992) that contributes significantly to the microlensing optical depth is consistent with the current data. Other populations (a thick LMC or material behind the LMC) will be even more difficult to constrain. We close by noting that no available data yet compel us to adopt either an intervening stellar or halo MACHO interpretation for all of the lenses. At least some lens events are associated with the Clouds themselves (MACHO-98-SMC-1 with high certainty, Rhie *et al.* 1999; MACHO-LMC-9 with less certainty, Bennett *et al.* 1996; Aubourg *et al.* 1999). Similar direct identification of even a few true halo lenses would have significant implications for the study of stellar evolution, galaxy formation, and baryonic dark matter.

ACKNOWLEDGMENTS: DZ acknowledges financial support from an NSF grant (AST-9619576), a NASA LTSA grant (NAG-5-3501), a David and Lucile Packard Foundation Fellowship and a Sloan Fellowship. DZ thanks Karl Gebhardt for the use of his maximum likelihood dispersion measurement code and Ann Zabludoff for

comments on drafts and discussions. Lastly, we thank David Alves, Jean-Philippe Beaulieu, Dave Bennett, and Carme Gallart for being gracious and providing helpful comments even though they disagree with the basic premise that the VRC stars are a significant lensing population.

References

- Alcock, C. *et al.* 1997a, *ApJ*, 486, 697
- Alcock, C. *et al.* 1997b, *ApJL*, 490, 59L
- Alcock, C. *et al.* 1998a, *ApJL*, 499, 9L
- Alcock, C. *et al.* 1998, *ApJ*, submitted (astro-ph/9807163)
- Alfonso, C. *et al.* 1998 in press
- Ansari, R. *et al.* 1996, *AA*, 314, 94
- Aubourg, E., *et al.* 1993, *Nature*, 365, 623
- Aubourg, E., *et al.* 1999, *AA*, in press (astro-ph/9901372)
- Beaulieu, J.-P., & Sackett, P.D. 1998, in press
- Bennett, D.P. 1998, *ApJL*, 493, L79
- Bennett, D.P. 1998, astro-ph/9808121
- Berterlli, G., Bressan, A., Chiosi, C., Fagotto, F., & Nasi, E. 1994, *A&AS*, 106, 275
- Binney, J., & Tremaine, S. 1987, *galactic Dynamics* (Princeton: Princeton Univ. Press)
- Bothun, G.D., & Thompson, I.B. 1988, *AJ*, 96, 877
- Castellani, V., Chieffi, A., & Pulone, L. 1991, *ApJS* 76, 911
- De Marchi, G., & Paresce, F. 1997, *ApJL*, 476, L19
- Dohm-Palmer, R.C. *et al.* 1997, *AJ*, 114, 2527
- de Vaucouleurs, G. 1957, *AJ*, 62, 69
- Efremov, Yu.N. 1978, *AZ*, 55, 272
- Evans, N.W., Gyuk, G., Turner, M.S., & Binney, J.J. 1997, *Nature*, in press
- Feast, M.W., Thackeray, A.D., & Wesseling, A.J., 1960, *M.N.R.A.S.*, 120, 64
- Feast, M.W., Thackeray, A.D., & Wesseling, A.J., 1960, *M.N.R.A.S.*, 121, 433
- Fitzpatrick, E.L. 1985, *ApJ*, 299, 219
- Fusi Pecci, F., Ferraro, F.R., Crocker, D.A., Rood, R.T., Buonanno, R. 1990, *AA*, 238, 95
- Gallart, C. 1998, *ApJ*, 495, 43
- Gardiner, L.T., & Noguchi, M. 1996, *M.N.R.A.S.*, 278, 191

- Geha, M. *et al.* 1998, *AJ*, 115, 1045
- Geisler, D., Bica, E., Dottori, H., Clariá, Piatti, A.E., & Santos, J.F.C. 1998, in *IAU 190 the Magellanic Clouds* (ed. Y.-H. Chu, N. Suntzeff, J. Hesser, & D. Bohlender)
- Gould, A. 1998, *ApJ*, 499, 728
- Graff, D. and the EROS 2 Collaboration 1998, in *IAU 190 the Magellanic Clouds* (ed. Y.-H. Chu, N. Suntzeff, J. Hesser, & D. Bohlender)
- Grebel, E.K. & Brandner, W. 1998, in *The Magellanic Clouds and Other Dwarf Galaxies*, eds. T Richtler & J. Braan, in press
- Harris, J., Zaritsky, D., & Thompson, I. 1997, *AJ*, 114, 1933
- Holtzman, J., *et al.* 1997, *AJ*, 113, 656
- Hughes, S.M.G., Wood, P.R., & Reid, N. 1991, *AJ*, 101, 1304
- Ibata, R.A., Lewis, G.F., & Beaulieu, J.-P. 1998, *ApJL*, submitted
- Iben, I., & Renzini, A. 1983, *ARA & A*, **2**, 271
- Johnston, K.V. 1998, *ApJ*, 495, 297
- King, C.R., Da Costa, G.S., & Demarque, P. 1985, *ApJ*, 299, 674
- Kent, S.M. 1986, *AJ*, 91, 1301
- Kim, S., Staveland-Smith, L., Dopita, M.A., Freeman, K.C., Sault, R.J., Kesteven, M.J., & McConnell, D. *et al.* 1998, *ApJ*, in press
- Kippenhahn, R., & Smith, L. 1969, *AA*, 1, 142
- Kunkel, W.E., Demers, S., Irwin, M.J., & Albert, L. 1997, *ApJL*, 488, L129
- Luks, Th., & Rohlfs, K. 1992, *AA*, 263, 41
- Majewski, S.R., Ostheimer, J.C., Kunkel, W.E., Johnston, K.V., & Patterson, R.J. 1998, in *IAU 190 the Magellanic Clouds* (ed. Y.-H. Chu, N. Suntzeff, J. Hesser, & D. Bohlender)
- Massey, P., Lang, C.C., DeGioia-Eastwood, K., & Garmany, C.D. 1995, *ApJ*, 438, 188
- Meatheringham, S.J., Dopita, M.A., Ford, H.C., & Webster, B.L. 1988, *ApJ*, 327, 651
- Paczyński, B. 1986, *ApJ*, 304, 1
- Payne-Gaposchkin, C.H. 1971, *Smithsonian Contr. Ap.*, No. 13
- Putnam, M. *et al.* 1998, *Nature*, in press
- Renault, C. *et al.* 1997, *AA*, 324, 69L
- Rhie, S.H. *et al.* , 1999, astro-ph/9812252
- Sahu, K.C. 1994, *PASP*, 106, 942
- Salpeter, E.E. 1955, *ApJ*, 121, 161

- Scalo, J.M. 1986, *Fund. Cosmic Phys.*, 11, 1
- Schild, R.E. 1977, *AJ*, 82, 337
- Schommer, R.A., Olszewski, E.W., Suntzeff, N.B., and Harris, H.C. 1992, *AJ*, 103, 447
- Shapley, H., 1956, *American Scientist*, 44, 73
- Shectman, S.A., Schechter, P.L., Oemler, A.A., Tucker, D., Kirshner, R.P., & Lin H. 1992, in *Clusters and Superclusters of Galaxies*, ed. A.C. Fabian (Dordrecht:Kluwer), 351
- Spitzer, L., & Schwarzschild, M. 1951, *ApJ*, 114, 385
- Spitzer, L., & Schwarzschild, M. 1953, *ApJ*, 118, 106
- Stetson, P.B. 1997, *Baltic Astron.*, 6, 3
- Udlaski, A., Kubiak, M., & Szymanski, M. 1997, *Acta. A.*, 47, 319
- Villumsen, J.V. 1985, *ApJ*, 290, 75
- Vogt, N.P. 1995, Ph.D. Dissertation, Cornell University
- Wielen, R. 1977, *AA*, 60, 263
- Zaritsky, D., Harris, J., & Thompson, I. 1997, *AJ*, 114, 1002
- Zaritsky, D. & Lin, D.N.C. 1997, *AJ*, 114, 2545
- Zhao, H. 1998a, *M.N.R.A.S.*, 294, 139
- Zhao, H. 1998b, *ApJL*, 500, 149




Fig. 1.— The stellar density plot (for $V < 21$) for the region in the LMC with reduced photometry from the Magellanic Cloud Photometric Survey. The central coordinates are roughly $\alpha = 5^h 20^m$ and $\delta = -66^\circ 48'$, the image is $\sim 4^\circ$ wide, with North at the top and East to the left. Each “pixel” corresponds to $30''$.




Fig. 2.— The final velocity template spectrum (constructed from the sum of the VRC stellar spectra).




Fig. 3.— Two tests of velocity uncertainties. The left panel shows the results of measuring cross correlation velocities for one calibration spectrum against the others. The distribution is Gaussian with low (3 km sec^{-1}) dispersion, demonstrating that the wavelength solutions are stable. The right panel shows the difference in the measured velocities with and without sky subtraction. The distribution is asymmetric but the dispersion of the fitted Gaussian is only 5 km sec^{-1} .




Fig. 4.— Repeatability of velocities. We show the results of comparing the radial velocity measured on one night vs. that measured on another night, normalized by the uncertainty estimate produced by the cross-correlation software (IRAF XCSAO).




Fig. 5.— The lower main sequences and a foreground population. We compare the CMD obtained by Holtzman *et al.* (1997) in the left panel to the same stellar population with the addition of a foreground population (uniformly distributed at a distance between 35 and 40 kpc along the line of sight) that is 8% of the LMC population. The lack of obvious differences between the two CMDs illustrates that even on the lower main sequence, where the statistics are best, it is difficult to exclude a foreground population.




Fig. 6.— The VRC/RC distribution. We plot histograms of vertical cuts taken through CMDs at the color of the red clump. The wing toward the left of the large peak is the VRC. The solid line are the data from ZL, the dashed lines represent results from three simulations. The adopted star formation history is listed above each corresponding panel.




Fig. 7.— The stellar density in $6' \times 6'$ pixels for selected main sequence stars ($19.15 < V < 19.5$) in the lower panel and the VRC in the upper panel. The total area shown in each panel is the equivalent to that shown in Figure 1.




Fig. 8.— The average ratio of the VRC stellar density to the main sequence stellar density as defined in the text as a function of main sequence stellar density (for $6' \times 6'$ boxes). The inset plot shows the entire data, the principal plot shows more detail and includes all but one of the data points. The solid curve represents the best fit model, the long-dashed curve represents a model with a fixed contribution of 14% of the MS stars to the VRC. The dashed horizontal line shows the predicted asymptotic value of the ratio for the Geha *et al.* star formation history.




Fig. 9.— The average ratio of the VRC stellar density to the main sequence stellar density as defined in the text as a function of main sequence stellar density (for $3' \times 3'$ boxes in filled circles and for $12' \times 12'$ boxes in open circles). The number of MS stars have been normalized to represent the numbers found per $6' \times 6'$ box. The solid curve represents the best fit derived from the data in Figure 8. The long-dashed curve represents the best fit model for the data from $3' \times 3'$ boxes. The dashed horizontal line shows the predicted asymptotic value of the ratio for the Geha *et al.* star formation history.




Fig. 10.— The VRC radial velocity distributions. In the left panel we show the radial velocities for all VRC candidates with a significant correlation value. Due to the nature of the velocity template resolution, the velocity is with respect to the mean of the large peak of VRC stars. In the right panel, we expand around that peak and overplot the best fit Gaussian model.




Fig. 11.— The RC radial velocity distributions. We show the radial velocities for all RC candidates with a significant correlation value. Due to the nature of the velocity template resolution, the velocity is with respect to the mean of the large peak of VRC stars (cf. Fig 10). The best fit Gaussian, convolved with the observational uncertainties, is shown as the solid line.




Fig. 12.— The radial velocity distributions of supergiants in the LMC (Feast, Thackeray, & Wesselink 1960,1961) after removing three stars within 1 kpc of the LMC center which had > 50 km sec^{-1} residuals from the mean rotation curve (see text). The dashed line shows the derived Gaussian for the entire population of stars (intrinsic velocity dispersion = 19.6 km sec^{-1}). The shaded bins indicate highly deviant stars that are suspected of distorting the derived Gaussian. The solid line shows the derived Gaussian after removing the three stars in the shaded bins (intrinsic velocity dispersion = 10 km sec^{-1}).